

THE ECOLOGICAL IMPORTANCE OF THE MARGHERITA DI SAVOIA SALTWORKS

CIRO ZENO

Atisale SpA
Environmental and Territorial Department
Via C. Colombo 112, 00147 Roma - Italy
e-mail: ciro.zeno@atisale.com

EXTENDED ABSTRACT

The Margherita di Savoia Saltworks, located in Apulia (South Italy), are the largest productive saltworks in Italy. They are connected with the Apulian wetlands, an important network thanks of its central geographic position, between the east and west of the Mediterranean basin. Several species and habitats of European and international interest (Natura 2000 network, Ramsar list) are present in them.

It plays a significant role as an area of stopover, wintering and breeding along the migratory routes of birds that cross the Mediterranean. The most interesting migratory and wintering species is the *Numenius tenuirostris* (slender-billed curlew), which is the bird most at risk of extinction in Europe.

Moreover in the early 1990s The *Phoenicopterus ruber* (greater flamingo) has colonised the reserve in great number, around 6,000 today, making it the largest concentration of the species in mainland Italy.

The Margherita di Savoia saltworks are entirely a man-made area, the characteristics of which (water levels, salinity) are preserved entirely due to sea salt production, which guarantees all the chemical and physical factors necessary for the survival of these habitats. It is also worth noting the type of production adopted in the saltworks, which makes it possible to recover the processing brine, thus eliminating one of the critical factors in the symbiotic relationship between salt production and environmental protection.

Key words: Lake Salpi, long-term salt production, "big beam" harvest, Apulian wetlands, greater flamingo Natura 2000 network, Environmental Management, Habitat Directive 92/43

1. THE MARGHERITA DI SAVOIA SALTWORKS

The Margherita di Savoia saltworks are the largest marine saltworks in Italy, with average annual production of approximately 5,500,000 quintals of salt, placing them amongst the top European saltworks in terms of volume and size.

They are located in the Basso Tavoliere area of Apulia (Coordinates: 41°24'N 016°04'E), a region in Southern Italy, immediately behind the Gargano promontory, inside the Gulf of Manfredonia. They take their name from the town of Margherita di Savoia.

The saltworks run lengthways, parallel to the Adriatic coast, for a distance of approximately 20 km, with a maximum width of 4 km. They cover a surface area of approximately 4000 hectares, with 3,781 subject to protection.

The Margherita di Savoia saltworks do not have a precise date of birth. There was originally a vast coastal lagoon on the site of today's saltworks. This lagoon was known as **lake Salpi** and was formed of brackish water and salt marshes. It gave its name to the

ancient Salapia, the Daunian lagoon city.¹ Following periodic episodes of sea water flooding the land behind the dunes, the evaporation formed extensive salt incrustations.² This was probably the event that caused man to become interested in this site, setting it aside for salt production. This activity has continued uninterrupted over the centuries, right up to today.



Figure 1: Margherita di Savoia saltworks

The Margherita di Savoia saltworks have always been eminently productive and have thus been a much sought after prize for the various peoples who have had control of them from the 6th century B.C., over the course of at least 2,500 years. As a result, the settlements that developed in the area have been destroyed and rebuilt many times over the centuries. The vicinity of lake Salpi also made the zone subject to malaria. Following major reclamation work in 1936 (Salpi Nuovo) and the 1950s (Alma Dannata), the entire lake was annexed to the salt works, considerably increasing their surface area. It was during this period that the theoretical study of Italian salt works reached its highest level of expression, developing a precise physical and mathematical model for the salt production process (*Iommi G*), which was used to remodel Italian saltworks and remained substantially unchanged until the mid-1980s. On the basis of this theory, the salt evaporation pond, which is seven times larger than the salt pan, should be subdivided into four zones of increasing saline density, until reaching the maximum density of 25.7°Be. At this point, the sodium chloride begins to precipitate and the water is poured into the salt pans.

Since the mid-1980s, following the economic crisis in the sector, a new method of deposited salt production has been implemented. This is known as **long-term**

¹ The ancient Daunians inhabited the northern part of Apulia, including the Gargano district from the 8th century B.C. onwards

² Undoubtedly, this area possesses all the factors that encourage the production of sea salt. First and foremost, the nature of the terrain: the salt works lie in a naturally impermeable basin comprised of silt banks with flaser bedding with an average impermeability coefficient of approximately 10^{-6} cm/s, which persists on clastic quaternary terrain of different compositions and granulometry.

The meteorological data is also undoubtedly favourable, with the average annual temperature being 16°C. In the period from May-October, the average maximum monthly temperature climbs to almost 23°C in May and 30-31°C in July and August, then falling to around 22°C in October. Thanks to the favourable thermal conditions and good ventilation, evaporation reaches very high values over the course of the year (1,665.4 mm), almost 70% of which occurs in the six months from May to October. Lastly, the pluviometric conditions are also extremely good and it should be pointed out that Apulia is home to the two areas with the lowest rainfall in all of mainland Italy.

production and is designed to make a significant reduction in production and operating costs.

During this process, the salt produced during the salt-extraction campaign (May - September) is not collected, but left deposited on the bottom of the salt pans. This process is repeated for 4-5 years. At the end of the cycle, there are several superimposed layers, with a total thickness that varies from 50 to 60 cm. In order to protect the layer of salt deposited on the bottom of the salt pans from atmospheric agents, the processing waters (those formed when the salt pans are emptied for harvesting) are used to form a protective cover of 30-40 cm thick.



Figure 2: Salt Harvest

The salt pans are harvested on a six-monthly basis using the **“big beam” harvest** method, which has led to noticeable simplification of the harvesting system. In fact, the thickness of the crust of salt deposited in the salt pans allows a certain number of diggers (3 or 4) to descend directly into the salt pan and form roads of salt (the so-called "big beams"). The thickness of these roads is double that of the crust. A scraper then compacts the “beams”, enabling the lorries to descend directly into the salt pan for loading by the diggers.


Several advantages have been made apparent by the adoption of long-term production and "beam" harvesting:

1. The harvest is separated from the production phase - something that is essential in the case of annual production. The harvest can therefore be scheduled on the basis of market requirements or quieter times of year at the plant, such as March and October.
2. The bottoms of the salt pans no longer require maintenance work, which involved compacting and pressing them so as to have completely flat bottoms.
3. The processing brine (bittern), which is normally disposed of at sea by salt works that implement annual harvests and is one of the critical factors in the relationship between sea salt production and the environment (*Davi J*), are instead reused to form a protective cover for the salt pans.
4. Less anthropogenic impact on the salt pans, inasmuch as the same area is harvested every five years.

5. Low anthropogenic impact as regards the harvesting method, due to the simplicity of the means and the small number of people involved: 4 diggers, one scraper, 10 lorries and 10 operators.

2. THE APULIAN WETLANDS

Apulia is home to numerous and extremely diverse coastal wetlands, which, environmentally speaking, are some of the most important in Italy. This system is formed of 38 wetland zones, covering a total area of approximately 24,000-26,000 hectares. In addition to providing extensive and varied habitats, it is also important because of its central geographic position, acting as a bridge between the east and west of the Mediterranean basin. It plays a significant role as an area of stopover, wintering and breeding along the migratory routes of birds that cross the Mediterranean. In particular, the large salt work complex is an irreplaceable stopping and nesting point for the thousands of migratory birds that visit the area in the winter months on their return from the Balkans, after crossing the Adriatic.



In order to analyse the importance and value of the salt works, we will use the Network Natura 2000 database for Apulia, created pursuant to Directive 79/409 Birds and Directive 92/43 Habitat.

Network Natura 2000 uses standardised procedures and criteria relative to the presence of habitats and rare and threatened species to identify a network of sites that are indispensable to the conservation of biodiversity in Europe. It is therefore a useful reference point for practical reports and the assessment of the environmental value of Apulia's wetlands.

We shall highlight their value and importance on the basis of the species and habitats of European interest (Annex I and II of EC Habitats Directive 92/43/EEC). The Apulian wetlands are known to provide habitat for the following priority species; fish, *Knipowitschia panizzai*, *Nono Aphanis fasciatus*; reptiles *Caretta caretta*, *Emys orbicularis*, mammals *Lutra lutra*; *Monachus m.*; thirty-two species of nesting birds, not to mention the hundreds of bird species that winter there, It provides twenty-six habitats, six of which are priority habitats (Sigismondi A.)

Some of the charts drawn up by the Region of Apulia Parks Department, regarding the region's wetlands, are featured below.

Table 1 - No. European species per wetland zone
Source: the Region of Apulia Parks Department





Table 2 - No. of exclusive species per site
Source: the Region of Apulia Parks Department




From these data we can extract these considerations:

In The Margherita di Savoia Salt works winter at least 10 species from Network Natura 2000, of which seven winter exclusively in this site.

This area has the minimum number of threat factor among the apulian wetlands, only four (A2,A5,A10,A11).

Table 3 - No. of threat and alteration factors per site
Source: the Region of Apulia Parks Department



Threat factors

A1: Reclamations; A2: Modifications and transformations of the habitat (constructions buildings, roads, ports, concrete fluvial embankments, climatic variations due to anthropic influences, obstructions on the water course, water exploitation); A3: Alteration of the water capacities; A4: Ground filling; A5: Regression of the coast, alteration of the dunale system; A6: increase of the salinity; A7: Water Pollution; A8: Reduction and impoverishment of the primary producers; A9: Agriculture; A10: Activity of the free time (nautical tourism, bathing, sports, sport fishing, birdwatching, climbing); A11: Hunting; A12: Illegal hunting and fishing; A13: Fish farming, problems of management, excessive fishing; A14: Introduction species alloctone;

3. THE “MARGHERITA DI SAVOIA SALTWORKS” WETLAND

The information collected on nesting birds over the last thirty years demonstrates the presence of numerous species of international and European interest in this site, some of which have population levels of national importance.³ Consequently, this area is subject to a series of restrictions. On the basis of Italian law, the National Forestry Service in the Gargano Area is responsible for protecting the area.

On 10th October 1977, the Ministry of Agriculture and Forestry included the salt works in the list of Italian State Protected Areas, classifying them as an Animal Nature Reserve.

On 30th May 1979, the Margherita di Savoia salt works were declared a wetland of international importance, pursuant to the Ramsar Convention (Iran, 1971),⁴ in virtue of the criteria of 1% of the population of the following species: *Tadorna tadorna* (sheldrake), widgeon, *Recurvirostra avocetta* (avocet) (criteria 6) and with the criteria of 20,000 birds (criteria 5)⁵.

³ the national importance is confirmed by the fact that the area is home to 1% of the national wintering population of 21 species of waterbirds.

⁴ “...This is one of the most important wetland zones in the Mediterranean and the most important in central and southern mainland Italy. It is a representative example of a water-dependent community characteristic of its biogeographical region. It represents one of the largest wetland areas in the Southern Adriatic that does not run the immediate risk of environmental transformation. It is located in a position that means that it can be readily used for scientific research and education. It is free from risks of major consequences as a result of external pollution, hydrobiological interference, alternative use of the territory and industrial activities ...”.

⁵ From the Ramsar Convention criteria for Identifying Wetlands of International Importance

Criterion 5: A wetland should be considered internationally important if it regularly supports 20,000 or more waterbirds.

Criterion 6: A wetland should be considered internationally important if it regularly supports 1% of the individuals in a population of one species or subspecies of waterbird.

They are included in the Natura 2000 network, as site ZPS IT9110006 "Saline di Margherita di Savoia" of 4,860 hectares, falling almost entirely within pSIC IT9110005 "Zone umide di Capitanata" of 14,109 hectares". Moreover, they belong to the Italian IBA network - area Important Birds – IBA: 203 *Promontorio del Gargano e zone umide della Capitanata*, promoted and coordinated by *BirdLife International* (www.birdlife.org.) whose aim is to identify, monitor and protect key sites for birds all through joint efforts of staff and volunteers at local, national and international level.

The progressive increase in salinity and the varying depths of the salt pans, the presence of banks and channels, alternating with expanses of mud and the abundance of phytoplankton and zooplankton biomass, create a series of ecological niches that are ideal habitats for different species of aquatic birds. The birds occupy these areas on a seasonal basis, in keeping with their needs.

Therefore, by visiting the various areas of the salt works, it is possible to observe, within just a few kilometres from each other, aquatic species such as ducks and coots, typical of wetlands with relatively deep, non-salty water, and greater flamingos, which prefer the saltiest ponds, where the chemical and physical conditions of the waters would be prohibitive for other species.

The wealth of fauna in the area can also be explained in terms of the considerable dimensions of the wetlands, with their geographical position on migratory routes and their interaction with other Apulian wetlands. These salt works are therefore a fundamental stopping point during the migration of birds from breeding sites in Northern Europe to African wintering quarters.

The National Wildlife Institute annually makes the census of the wintering and nesting birds inside the Saltworks and periodically publishes the results. The number of wintering birds varies greatly from one year to the next, primarily depending on the weather conditions. It oscillates between 20 and 50,000 (a peak of 58,709 birds was recorded in 2000).

Hundreds of different bird species have been observed in the salt works. There also many other species of European interest, with some of the most regular visitors being the *Phalacrocorax carbo* (Pygmy Cormorant), the *Egretta alba* (Great white Egret), the *Egretta garzetta* (Little Egret), the *Botaurus stellaris* (Great Bittern), the *Ciconia ciconia* (White Stork), the *Ciconia nigra* (Black Stork), the *Platalea leucorodia* (Spoonbill), the *Phoenicopterus ruber* (Greater Flamingo), the *Aythya nyroca* (Ferruginous Duck), the *Pluvialis apricaria* (Golden Plover), the *Philomachus pugnax* (Ruff), the *Tringa glareola* (Wood Sandpiper), the *Limosa lapponica* (Bar-tailed Godwit), the *Phalaropus lobatus* (Red-necked Phalarope), the *Hydroprogne caspia* (Caspian Tern), the *Sterna sandvicensis* (Sandwich Tern), the *Chlidonias niger* (Black Tern), the *Falco biarmicus* (Lanner Falcon), the *Falco peregrinus* (Peregrine Falcon), the *Circus aërginosus* (Marsh harrier), the *Circus cyaneus* (Northern harrier), the *Pandion haliaetus* (Osprey) and the *Asio flammeus* (Short-eared Owl).

The sea beds are home to large populations of bivalves, which are a source of food for many bird species such as the sheldrake.

The reserve is also an important breeding ground for birds. It is home to the largest Italian colonies of *Recurvirostra avocetta* (Avocets), with over 650 pairs, of *Larus melanocephalus* (Mediterranean Gulls), with approximately 1000 pairs in some years, of *Larus genei* (Slender-billed Gulls), with sometimes over 900 pairs being recorded, of *Sterna albifrons* (Little Terns), with up to 600 pairs, and of *Himantopus himantopus* (Black-winged Stilts), with up to 140 pairs .

But the most interesting migratory and wintering species, from a conservationist point of view, is the *Numenius tenuirostris* (slender-billed curlew), which is the bird most at risk of extinction in Europe. The Gulf of Manfredonia is designed by the National Action Plan like the most important target area for the action of preserving this bird from the risk of

extinction, because it provide the most important stopping point for this bird on a national level and one of the most important in Western Europe⁶.





Figure 3. Avocets

Some nests in the salt works are important not just in terms of numbers, but also in terms of the rarity of the species, such as those of the *Gelochelidon nilotica* (gull-billed tern), the *Sterna sandvicensis* (sandwich tern) and the *Larus melanocephalus* (Mediterranean gull).

Table 4: Increase in birds nesting in the Saltworks
Source: the Region of Apulia Parks Department




The *Phoenicopterus ruber* (**greater flamingo**) represents a special case, inasmuch as this species has colonised the reserve in great numbers over recent years. Flamingos began to settle in the shallower, saltiest ponds in the early 1990s, managing to breed within the salt works for the first time in 1996. The number of individuals has thus

⁶ Slender-billed Curlew is present in the maximum risk categories identified by the main international conventions and community directives to faunal subject. It is enclosed I the Appendix I of the of the CEE 79/409, in the Appendix I of CITES, In the Appendix I of the Convention of Bonn (Conservation of Migratory Species of Wild Animals) and in the Annex II of the Convention of Bern (Conservation of European Wildlife and Natural Habitats). The Convention on Migratory Species of Wild Animals (CMS) produced, in 1994, a suitable memorandum book of Understanding concerning conservation measures for the Slender-billed Curlew (MoU), subscribed by several other countries included in the list of the Ranges States, among which Italy (signature affixed on April 18th, 2000). In this document the kind is expressly declared to extinction risk, act what to the senses of the Italian law no. 157/92, article. 2, paragraph 1, c letter, allows to attribute her the status of kind especially protected.

continued to climb, until reaching around 6,000 today, making it the largest concentration of the species in mainland Italy. These birds have therefore become a stable presence in the protected area, where their food abounds, primarily in the form of the crustacean *Artemia salina*, the only invertebrate that can live in water with salinity of over 300‰, to they owe their pink colouring.

Table 5: Flamingo population trends
Source: the Region of Apulia Parks Department



4. CONCLUSIONS

Among the Apulian wetlands, the saltworks of Margherita di Savoia is certainly the one has the better conditions for the conservation of its ecosystem and guarantees the safeguard of the biodiversity. (tab. 3). It's importance in the national and international wetland context is well recognized. On the other these habitats have elements of fragility due to the difficult interaction with the surrounding territory and to the lack of boundary buffer areas. For this it's necessary to achieve a greater involvement of the interested municipalities, in order to develop a shared model of management of the territory, aware of the needs of environment, and in which it's clearly recognized the irreplaceable role of the sea salt production. Probably this is a common situation for other Saltworks.

From this perspective some actions can be employed by the private company that own the Saltworks:

- 1) Implement a system of environmental management, like ISO 14001 and, for the European saltworks, EMAS standards.
- 2) Promote scientific studies who, starting from the salt production, examine in exhaustive and consistent way all the connected aspects: naturalistic, biological, geologists and microbiologic.
- 3) If the Saltworks belong to European Natura 2000 network, promote the adoption of the Management Plan (article. 6 EC Habitats Directive 92/43/EEC).

REFERENCES

1. Iommi G. (1941), *Le Saline del Monopolio Italiano nell'Anno XVIII dell'Era Fascista, The saltworks of Italian Monopolies in the XVIII of the Fascist Age*. Istituto poligrafico dello stato, Roma
2. Sigismondi A. Apulia Parks Department (2005), *Le saline di Margherita di Savoia, valore ed importanza nel sistema delle zone umide della Puglia. Atti del convegno, La produzione del sale e la salvaguardia dell'ecosistema*. www.atisale.com
The Margherita di Savoia Saltworks: value and importance in the Apulian wetlands networks, *Proceedings of the Conference*, The production of sea salt and the safeguard of the ecosystem, www.atisale.com
3. Zeno C. – Sereno F. (2002) Margherita di Savoia: the Queen of the Mediterranean saltworks. *Edizioni Cedel*, Margherita di Savoia
4. Panella M. (1998), *Information Sheet on Ramsar Wetlands: Saline di Margherita di Savoia*, <http://www.wetlands.org/reports/infosheet.cfm?siteref=3IT026>
5. Serra L., Magnani A., Dall'Antonia P., Baccetti N., (1997), Risultati dei censimenti degli uccelli acquatici svernanti in Italia (Census of wintering water-birds in Italy), *Biol. Cons. Fauna* 101
6. Joseph S. Davis (1999), Solar Saltworks An Environmentally Friendly Industry, *Proceedings of Post Conference Symposium Saltworks: Preserving Saline Coastal Ecosystems*. http://www.gnest.org/Conferences/Saltworks_post/031-037.pdf
7. Bissanti A. (1977), *Le saline di Margherita di Savoia: localizzazione e cambiamenti nell'organizzazione del territorio. La terra del sale, (Margherita di Savoia saltworks: localization and changes in the territory organization. The land of salt*
8. Istituto Nazionale per la fauna selvatica Risultati del censimento degli uccelli acquatici svernanti in Italia: distribuzione, stima e trend della popolazione in 1991- 2000 – ediz. Baccetti
National wildlife Institute - Results of the censuses of the wintering aquatic birds in Italy: distribution, estimate and trend of the populations in 1991-2000", edits in 2002 Baccetti
9. Zenatello M. e N. Baccetti (a cura di), 2001 - Piano d'azione nazionale per il Chiurlottello (Numenius tenuirostris). the Quad. Cons. Natura, 7, Min. Ambiente - Ist. Naz. Fauna Selvatica National Action Plan for Slender-billed Curlew (Numenius tenuirostris). Wildlife Min. *National wildlife Institute*
10. Ariel Brunner, Claudio Celada, Patrizia Rossi, Marco Gustin Sviluppo di un sistema nazionale delle ZPS sulla base della rete delle IBA (Important Bird Areas)" LIPU- BirdLife Italia www.lipu.it Development Of a ZPS national system under the IBA basis

. Slender-billed Curlew is present in the maximum risk categories identified by the main international conventions and community directives to faunal subject. It is enclosed I the Appendix I of the of the CEE 79/409, in the Appendix I of CITES, In the Appendix I of the Convention of Bonn (Conservation of Migratory Species of Wild Animals) and in the Annex II of the Convention of Berna (Conservation of European Wildlife and Natural Habitats). The Convention on Migratory Species of Wild Animals (CMS) produced, in 1994, a suitable memorandum book of Understanding concerning conservation measures for the Slender-billed Curlew (MoU), subscribed by several other countries included in the list of the Ranges States, among which Italy (signature affixed on April 18th, 2000). In this document the kind is expressly declared to extinction risk, act what to the senses of the Italian law no. 157/92, article. 2, paragraph 1, c letter, allows to attribute her the status of kind especially protected.

Il Slender-billed Curlew è inserito nelle categorie di massimo rischio individuate dalle principali convenzioni internazionali e direttive comunitarie a soggetto faunistico. È incluso nell'Allegato I della Direttiva CEE 79/409, nell'Allegato I della CITES, nell'Allegato I della Convenzione di Bonn e nell'Allegato II della Convenzione di Berna. La Convention on Migratory Species of Wild Animals (CMS) ha prodotto, nel 1994, un apposito Memorandum of Understanding concerning conservation measures for the Slender-billed Curlew (MoU), sottoscritto da numerosi altri paesi inclusi nella lista dei Range States, tra cui l'Italia (firma apposta il 18 aprile 2000). In questo documento la specie viene espressamente dichiarata a rischio di estinzione, atto che ai sensi della Legge n. 157/92, art. 2, comma 1, lettera c, consente di attribuirle lo status di specie particolarmente protetta.

Conseguentemente L'Italia ha elaborato un piano d'azione interessante tutto il territorio nazione con il quale ha individuato cinque